RESEARCH ARTICLE



Public perceptions of the COVID-19 pandemic management in Bangladesh: a qualitative exploration [version 1; peer review: 2 approved with reservations]

Taufique Joarder 1, Muhammad N.B. Khaled 2, Mohammad A.I. Joarder³

¹Public Health Foundation, Bangladesh, Dhaka, Bangladesh ²International Food Policy Research Institute, Dhaka, Bangladesh ³Kumudini Women's Medical College Hospital, Tangail, Bangladesh

 First published: 02 Mar 2021, 10:170 https://doi.org/10.12688/f1000research.28333.1
Latest published: 02 Mar 2021, 10:170 https://doi.org/10.12688/f1000research.28333.1

Abstract

Background: Since the emergence of the COVID-19 outbreak, Government of Bangladesh (GoB) has taken various measures to restrict virus transmission and inform the people of the situation. However, the success of such measures largely depends on a positive public perception of the government's ability to act decisively and the transparency of its communication. We explored public perceptions of pandemic management efforts by the Bangladeshi health sector decision-makers in this study.

Methods: As this qualitative research was conducted during the COVID-19 pandemic, data was gathered through seven online mixed-gender focus group discussions involving 50 purposively selected clinicians and non-clinicians.

Results: The study participants concurred that, from the outset, decision-makers failed to engage the right kind of experts, which resulted in poor pandemic management that included imposing lockdown in periphery areas without arranging patient transport to the center, declaring certain hospitals as COVID-19 dedicated without preparing the facilities or the staff, and engaging private hospitals in care without allowing them to test the patients for COVID-19 infection. Several participants also commented on ineffective actions on behalf of the GoB, such as imposing home guarantine instead of institutional, corruption, miscommunication, and inadequate private sector regulation. The perception of the people regarding service providers is that they lacked responsiveness in providing treatment, with some doctors misleading the public by sharing misinformation. Service providers, on the other hand, observed that decision-makers failed to provide them with proper training, personal protective equipment, and workplace security, which has resulted in a high number of deaths among medical staff.

Conclusions: The Bangladeshi health sector decision-makers should



Open Peer Review

- 1. Theresa Hoke, FHI 360, Durham, USA
- 2. **Manuela Villar Uribe**, World Bank Group, Washington, USA

Any reports and responses or comments on the article can be found at the end of the article.

learn from their mistakes to prevent further unnecessary loss of life and long-term economic downturn. They should adopt a sciencebased response to the COVID-19 pandemic in the short term while striving to develop a more resilient health system in the long run.

Keywords

COVID-19, Pandemic, Resilience, Health policy and systems research, Bangladesh



This article is included in the Disease Outbreaks

gateway.



This article is included in the Coronavirus

collection.

Corresponding author: Taufique Joarder (taufiquejoarder@gmail.com)

Author roles: Joarder T: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation; Khaled MNB: Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Writing – Original Draft Preparation; Joarder MAI: Formal Analysis, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Joarder T *et al*. This is an open access article distributed under the terms of the Creative Commons Attribution License , which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Joarder T, Khaled MNB and Joarder MAI. Public perceptions of the COVID-19 pandemic management in Bangladesh: a qualitative exploration [version 1; peer review: 2 approved with reservations] F1000Research 2021, 10:170 https://doi.org/10.12688/f1000research.28333.1

First published: 02 Mar 2021, 10:170 https://doi.org/10.12688/f1000research.28333.1

Introduction

In December 2019, an unknown pneumonia-like disease appeared in Wuhan, China, but rapidly spread across the globe, prompting the World Health Organization (WHO) to label it as Coronavirus Disease 2019 (COVID-19)1. On 30 January 2020, the WHO declared a Public Health Emergency of International Concern (PHEIC) followed by pandemic declaration on 11 March 2020². In Bangladesh, the first case of COVID-19 infection was detected on 8 March, resulting in closure of educational institutions on 16 March. Following the first death on 18 March, in an attempt to contain the spread of the disease, the Government of Bangladesh (GoB) declared a 'general holiday' from 26 March to 4 April, which was repeatedly extended until 9 April, 14 April, 25 April, 5 May, 16 May, and 30 May. Despite these measures, COVID-19 infections continued to increase, but lockdown (which was formally termed by the government as 'general holidays') was withdrawn on 31 May³. The number of confirmed COVID-19 cases in Bangladesh exceeded 100,000 on 18 June, and the upward trend continued throughout summer, with 200,000 cases recorded on 18 July, 300,000 on 26 August, and 400,000 on 27 October⁴. By November 2020, Bangladesh ranked 24th and 20th in the world with respect to the total number of cases and deaths, respectively. At one point, Bangladesh ranked 3rd in terms of the number of new cases per day, but given that the country's test rate (15,863 per million inhabitants) is among the lowest in the world (it is the second-lowest after Afghanistan among its South Asian neighbors) these figures are likely to be much higher⁵.

Pandemic response in Bangladesh is guided by the Infectious Diseases (Prevention, Control, and Elimination) Act 2018, which places the Directorate General of Health Services (DGHS) as

the central coordinating and responsible body for COVID-19 response. Institution of Epidemiology, Disease Control, and Research (IEDCR) is the main scientific body to provide technical guidance and support for screening at the point of entry, and is in charge of imposing quarantine, managing contact tracing, and conducting initial testing (which was later contracted out to other government and a few private laboratories), while also providing forecasting and surveillance services, and the overall outbreak response⁶. This agency was highly criticized for monopolizing all the COVID-19 tests in the first three weeks following the detection of the first case in a country of 180 million inhabitants. It was also blamed by the medical community for severe shortages of personal protective equipment (PPE) and tests, due to which many health professionals refused to provide care to infected individuals7. To address this issue, from 3 April, approvals for additional test facilities were gradually issued, initially in the public and later in private facilities, totaling to 117 on 18 November⁸. However, even though GeneXpert equipment was already available for tuberculosis test, this antigen-based rapid test was only made available for COVID-19 test by the government in July 20209. Major events related to the COVID-19 pandemic in Bangladesh are shown in Figure 1.

Since the COVID-19 outbreak, the GoB has taken various measures to inform the public of the situation and restrict the transmission of the infection. However, available evidence indicates that success of such measures largely depends on a positive public perception of government's ability to manage pandemics effectively, as well as foster multi-stakeholder cooperation¹⁰, garner social order within the population^{10,11} and ensure good governance^{12,13}. In this context, timely and transparent



Figure 1. COVID-19 timeline in Bangladesh.

communication is essential^{14,15}, as is involvement of technical and health experts in decision-making^{10,16}. Gathering and sharing information on newly infected individuals and their contacts (as a part of contact tracing activities) is an important pandemic response activity, which can only be effective if the general public trusts the relevant agencies and service providers^{10,14,16}. When people have a positive perception of the health system, they are more likely to adhere to any measures imposed to protect public health^{11,16,17}. Thus, COVID-19 response requires adaptive leadership capable of making bold decisions and passing timely regulations based on the most recent scientific evidence, which is impossible without a positive perception of or trust on decision-makers and all pertinent stakeholders, including general public¹⁸.

Although the epidemiologic features of SARS-COV-2 virus^{19,20}, its clinical manifestations in different patient groups^{21,22}, and its molecular characteristics²³⁻²⁵, as well as health systems response²⁶, economic and social consequences²⁷⁻²⁹, and public attitudes toward the measures implemented in Bangladesh have been investigated³⁰⁻³², public perceptions of pandemic management efforts by the responsible bodies have never been studied. Motivated by the work of Bigdeli et al., we decided to explore the public perceptions of COVID-19 pandemic management in Bangladesh by focusing on the relationships between (1) people and the decision-makers (or the larger health system governance), (2) people and the service providers (only physicians were covered in this study), and (3) service providers and decisionmakers³³. Findings yielded by this qualitative study will help decision-makers in introducing new or revising existing measures to allow service providers to better respond to the pandemic and increase public trust in the health system.

Methods

Study design

To gather the data for this qualitative study, seven focus group discussions (FGDs) were conducted as a part of which participants' perceptions of the COVID-19 pandemic management in Bangladesh were explored.

Ethical considerations

All the respondents provided verbal informed consent. All ethical principles were adhered to. The research was reviewed and approved by the Ethical Review Committee of the Public Health Foundation, Bangladesh (Reference number: 02/2020). During the pandemic, it was nearly impossible to gather all the participants of the FGDs and collect written consent. One may argue that it might be collected through digital signature. However, not all the participants were technologically well-equipped and trained. For convenience and treating all the participants' consent in the same manner, we collected verbal consent. The Ethical Review Committee (ERC) approved this procedure.

Study site, study population and participant selection

The study was promoted via a Google Forms link circulated across social media and email databases. First, the Google Forms link was circulated on 19 May among the participants of a webinar on health system trust, organized by a youth organization, the United Nations Youth and Students Association of Bangladesh (UNYSAB). The link was further circulated across social media and email databases, requesting expression of interest to participate in the study. The email databases of the members of the Public Health Foundation Bangladesh and the UNYSAB have been used in this regard. For circulating the form, the link along with a request to fill-up the form, was posted in the network of social media groups of researchers, health professionals, and university-based organizations.

Then, from the list of all the interested persons, the participants were purposively selected such that the FGD participants could be broadly classified into clinicians (graduate students with medical or dental background pursuing degrees in public health at a private university; renowned public health experts with a medical background; and clinicians practicing medicine or dentistry) and non-clinicians (undergraduate students pursuing non-medical degrees such as management, marketing, botany, business, and pharmacy, etc. at a public university; undergraduate students pursuing public health degrees at a public university; undergraduate students pursuing degrees in food and nutrition at a public university; and different professionals such as executives, trainers, managers, and coordinators of public and private organizations).

Data collection

Prior to commencing the FGDs, we developed a discussion guide in Bangla (see Extended data³⁴), which was pilot tested in a session where a large group of health professionals and university students were present, organized by the UNYSAB, and adjusted in terms of issues addressed and the pattern of the language, where necessary. The goal was to focus discussions on the topics pertinent to this investigation, i.e., participants' perceptions of COVID-19 pandemic management by the health sector decision-makers and the service providers, the implications of the actions (or lack thereof) taken by the decision-makers and service providers, and suggestions for improvements in pandemic management strategies.

Once a sufficient number of participants of both genders was recruited, seven FGDs (which were recorded through the video conferencing software Google Meet) were conducted between 15 and 17 June 2020, each involving 6–10 participants.

The FGDs were moderated by the first author (male), who has a doctorate in public health and is a health policy and systems researcher with experience and expertise in qualitative research methods. As a researcher in public health, he knew the participants in the FGD's with the health professionals and these participants knew him as well. The second author (male), trained in economics with a Master's degree and experienced in qualitative research, assisted in notetaking. He, however, had no prior engagement with any of the participants. Each FGD lasted 60–105 minutes and was conducted in Bangla, the native language of the respondents and the researchers. The research interests were explicitly explained to the participants, but the floor was open and no leading discussion points were initiated.

Data analysis

Prior to content analysis-chosen due to the scarcity of existing literature on pandemic management in Bangladesh-the FGDs were transcribed by the research team³⁵. For the diversity of the role among the FGD participants working in the health sector, data and observations were unique. In the FGD's conducted among the university students, data saturation took place. Thematic analysis commenced with listening to the recordings and reading the transcripts, which allowed a coding schema to be developed in Microsoft Excel (version: Professional Plus 2016) based on the questions asked, noting the first impressions followed by labelling the text segments by newly emerging codes. Next, similar-meaning codes were merged and sorted into broader categories. To substantiate the emerging themes, appropriate excerpts from the FGDs were identified. In order to increase validity, the first and second author independently coded the dataset, seeking input from the third author in case of any disagreement.

Results

Background characteristics of the focus group participants

In total, 50 individuals (28 males and 22 females, aged 19–75 years) took part in seven FGDs (Table 1). Four of these FGDs were held with individuals with a non-clinical background (n = 28) and the remaining three capture the views of clinicians (n = 22). Nearly half of the respondents had training in public health.

Participants' perceptions of the health system decisionmakers

Participants' perceptions regarding health systems decisionmakers have been presented under three sub-themes: their perceptions regarding the preparatory phase of the pandemic (not appointing the appropriate professionals, leading to incorrect management steps), coordination (indecisions stemming from incoordination), and actions by the health sector decisionmakers (miscommunication, poor regulation).

Perceptions regarding preparation

For years, the Bangladeshi health system has been undermined by budget shortages, lack of quality services, high out-ofpocket payments, unregulated private sector, and a highly centralized secondary or tertiary care. These issues hindered the pandemic response, as the right persons were not placed in the right positions at the outset, as explained by a professor of public health:

"An epidemic is a public health emergency; it is neither clinical nor an administrative issue. So, we must see this problem through the public health lens. We [epidemiologists] already know what to do to control an epidemic. ... We the public health professionals should be given the flexibility that we are free to do whatever is needed for the country, not something that just pleases the political leadership." [FGD-6, renowned public health experts, clinical background]

FGD number	Group characteristics	Number of respondents	Age range in years	Male/ female	Clinical or non- clinical background
FGD-1	Undergraduate students pursuing degrees in management, marketing, botany, business, and pharmacy at a public university	6	19–21	5/1	Non-clinical
FGD-2	Graduate students with medicine or dentistry as their undergraduate background pursuing public health degrees at a private university	6	25–34	0/6	Clinical
FGD-3	Undergraduate students pursuing public health degrees at a public university	9	21–26	4/5	Non-clinical
FGD-4	Undergraduate students pursuing food and nutrition degrees at a public university	6	22–25	0/6	Non-clinical
FGD-5	Different professionals such as executives, trainers, managers, and coordinators of public and private organizations	7	24–28	6/1	Non-clinical
FGD-6	Renowned public health experts with medical background	7	45–75	5/2	Clinical
FGD-7	Practicing clinicians with either medical or dentistry background	9	28–67	8/1	Clinical

Table 1. Characteristics of the focus group discussion (FGD) participants.

Failure to engage the right professionals in the decisionmaking resulted in a rapidly escalating public health crisis, as the testing capacity, medical equipment and PPE provisions in the health centers were lacking. Study participants also concurred that the crisis was exacerbated by not instituting subsistence allowance for the poor before imposing lockdown (denoted by the GoB as 'general holiday'), by failing to allow sufficient time for families to prepare for shop closures, and for increasing uncertainty by extending lockdown on a weekly basis. In addition, the health system actors wasted critical time by initially conducting tests at a single government facility before allowing private centers, although few in number, to engage in testing and provide COVID-19 care. Non-government organizations, and National Tuberculosis Control Program of the government had access to antigen-based rapid test, GeneXpert machines, which were utilized at a much later stage of the pandemic. As one student of public health explained:

"Government is allowing, although lately, private diagnostic centers to perform COVID-19 tests. But there are many other machines which are not yet being utilized." [FGD-2, graduate students of public health at a private university, clinical background]

Perceptions regarding coordination

Several FGD participants commented on lack of coordination which manifested through imposing lockdown in periphery areas of the country without arranging for the patients to be transported to the centrally located health facilities, declaring several hospitals as COVID-19 dedicated institutions without providing the required resources, and involving private hospitals in care provision without allowing them to test patients for COVID-19 on admission. Additionally, they were of view that lockdown was terminated due to the upcoming major Muslim religious holiday—the Eid—even though the outbreak was not showing any signs of abating. This decision was reached despite objections from the leading public health experts. Reflecting on these issues, a renowned public health expert remarked:

"The civil administration must coordinate with the health system people at district and sub-district levels to ramp up the response against COVID-19. Success cannot be achieved without a tactful decentralization, involving different relevant ministries." [FGD-6, renowned public health experts, clinical background]

According to one FGD participant, lack of coordination contributed to the rapid spread of the disease among the low-income garment employees of Bangladesh:

"BGMEA [national trade organization of garment manufacturers] ordered garment workers to return to Dhaka to save their jobs. After they returned by thousands, braving the coronavirus infection, BGMEA declared that the factories will not open. They [BGMEA] never consulted public health experts or health department. On the other hand, the police announced that they will not allow anyone to enter Dhaka. The innocent workers were caught between a rock and a hard place due to the lack of coordination between different departments." [FGD-2, graduate students of public health at a private university, clinical background]

Perceptions regarding actions taken by the health sector decision-makers

Several actions by the health sector decision-makers were openly criticized by the study participants. They were particularly critical of the decision to impose home rather than institutional quarantine at the beginning of the pandemic, even though intimate Bangladeshi culture is not conducive to home quarantine. They were also of view that point-of-entry screening was weak, and blamed widespread corruption for PPE shortages, purchases of sub-standard equipment and mismanagement of relief materials. These issues, along with miscommunication, were frequently discussed in media. Several participants also criticized the decision to disguise lockdown as a 'general holiday' in order to reduce panic, while failing to allow the residents enough time to prepare. This was aptly surmised by one participant, who noted:

"When they [government] say it is a 'general holiday' instead of 'lockdown', people confuse it with something festive. That's why we saw people going to Cox's Bazar [a popular tourist destination] for tourism purposes. Some of my friends even got married during this period, taking advantage of the 'general holiday'." [FGD-3, undergraduate students of public health at a public university, non-clinical background]

Poor regulation was another complaint voiced by many participants, who were of view that it caused rapid escalation in prices of essential goods due to panic buying, while permitting uncontrolled advertisement and sales of unproven COVID-19 medicines (e.g., hydroxychloroquine, ivermectin, remdesivir, etc.). One participant commented on the proliferation of unauthorized and even fake testing centers that were providing false COVID-19-negative certifications:

"Today, I saw in the news that someone who had been found corona negative here was found positive after landing in Japan. So, when things like these happen, our trust is compromised. These instances may even adversely affect our foreign relations." [FGD-5, service holders of different professions, non-clinical background]

Participants' perceptions of the service providers

Participants' perceptions regarding health service providers include lack of responsiveness of the providers, spreading misinformation, colluding in corruption.

When the pandemic started, some newspapers alleged that, fearing for their own safety, some doctors were refusing to provide service to COVID-19 patents, or were not responsive enough while providing treatment. As one public university student explained:

"In hospitals, especially the government hospitals, doctors don't care about the patients. Doctors should not only provide clinical care, but also explain the disease, talk to the patient with respect, and provide more time." [FGD-1, undergraduate students of different departments at a public university, non-clinical background]

Some doctors were also accused of spreading misinformation through social and mainstream media. In a video that was rapidly disseminated across social media, one doctor confidently claimed that coronavirus would go away in the summer, while another respected senior doctor openly advertised unproven medicines on TV. These actions were condemned by a participant:

"I found many of my doctor friends posting about different treatments for COVID-19. I think this may confuse and mislead people, as different doctors are saying different things." [FGD-2, graduate students of public health at a private university, clinical background]

Since the COVID-19 pandemic began, a doctor has been found colluding with someone guilty of running unauthorized testing centers³⁶. Some were found promoting unproven medicines, and providing false certificates of COVID-19 negativity. One medicine (hydroxychloroquine) was initially included into the national treatment guidelines, only to be subsequently removed following the WHO's warning about its ineffectiveness. A public health expert attributed these decisions to the vested interest of some clinicians serving on the technical committee in promoting certain treatments:

"Some [doctors] are saying plasma therapy is the solution, some are promoting different other drugs like hydroxychloroquine even though the WHO is saying there is no specific treatment for COVID-19." [FGD-6, renowned public health experts, clinical background]

Perceptions of the service providers about the decisionmakers

Service providers' perceptions regarding health systems decision-makers include leaving them unprepared and untrained in the face of the pandemic, not recognizing their sacrifices and the lack of workplace security stemming from COVID-19 mismanagement.

Doctors that took part in this investigation felt that the health system decision-makers failed to prepare them adequately to combat COVID-19 effectively. They complained about lack of training, absence of treatment guidelines, PPE and equipment shortages, as well as inadequate food and logistic support while on duty. On this, a doctor that has been on COVID-19 duty since the pandemic outbreak said:

"Doctors did not receive proper training or treatment guidelines, only online training, nothing on triage, how to handle indoor patients, no idea about treatment guideline, nothing on donning and doffing of PPE, or mental stress management. I feel like swimming in an unfathomable sea, without proper training." [FGD-7, practicing clinicians, clinical background]

These issues resulted in a very high number of deaths among Bangladeshi doctors, which further undermined healthcare providers' trust in the health system. Despite their sacrifices, they were not granted prioritized testing or healthcare, while also experiencing delays in salary payments. As a result, many doctors lacked motivation, as explained by one participant:

"Government did not clarify direction regarding who would get the motivation package. Some doctors did not even receive their regular salary. This demoralized the doctors. ... I know several young doctors who are saying that, if they are assigned COVID-19 duty, they will simply resign." [FGD-6, renowned public health experts, clinical background]

Several doctors expressed concerns over workplace security, as Bangladeshi people take out their dissatisfaction over the health system inadequacies on the doctors. To highlight the growing violence which resulted in a death of a colleague, many doctors stopped telemedicine services which they were previously offering benevolently to combat the COVID-19 crisis. A physician engaged in COVID-19 response said:

"Decision-makers should make the work environment of the doctors safe in such a way that they themselves would confidently send their own children for treatment. Many of them [health sector decision-makers] have children who are doctors. I personally know several of them who are forbidding their children to serve in COVID-19 units, because there is no security there." [FGD-7, practicing clinicians, clinical background]

Discussion

The findings yielded by this qualitative study indicate that several problems emerged as a consequence of failure to engage the right kind of experts in managing the pandemic. As a result of poor decision-making, the Bangladeshi health system was inadequately prepared to respond to the COVID-19 outbreak, as evident in the negative perception of our participants regarding the service providers especially in terms of quality of care they provided, and misinformation some of them shared in the social and mainstream media. Service providers also complained about lack of training, PPE, equipment, motivational packages, and workplace security.

The finding that the Bangladeshi health system failed to engage the right experts in the right positions is supported by several news articles and reports covering this topic. The Government of Bangladesh formed a 17-member National Technical Advisory Committee (NTAC) on 19 April 2020, more than a month after the first COVID-19 case was detected in the country. In the interim, most of the pandemic control efforts were entrusted to bureaucrats or administrators, many of whom lacked expertise or experience in health, let alone pandemic management. It is also worth noting that only three members of the NTAC had a public health career track³⁷. This issue was further compounded on 21 April, when the government assigned 64 top bureaucrats to supervise and coordinate relief distribution activities in 64 districts of Bangladesh³⁸ without seeking input or technical leadership from public health professionals. A policy analysis on the human resources for health in Bangladesh revealed that the DGHS is principally managed by

the clinicians at the expense of public health experts. The same applies to the Ministry of Health and Family Welfare level, which comprises of members drawn from other ministries often unrelated to the health sector³⁹. Given that such administrative approach is not conductive to pandemic management, lessons can be learned from Switzerland, Georgia, and New Zealand and other countries where science-based public health strategies have been proven highly effective⁴⁰.

Since doctors are often seen as the face of a health system, people blame them for any inadequacies in care delivery despite considerable sacrifices most doctors have made throughout the pandemic. So far, around 3,000 doctors in Bangladesh have contracted the virus and more than 100 have died due to COVID-19⁴¹. The negative perceptions regarding the service providers have been widely reported in Bangladeshi media⁴², which were attributed to poor communication skills and inadequate responsiveness (i.e., addressing the social needs of the patients such as being treated with friendliness, respect, information, trust, and sensitivity) in recent academic studies⁴³⁻⁴⁵.

Service providers' claims that inadequate training, PPE and equipment shortages are the main cause of their grievances have also been documented in other studies from Bangladesh. In a study conducted from 9 to 14 April 2020, Islam and colleagues examined the frontline health workers' perceptions and opinions on their personal safety while attending COVID-19 patients. Their findings show that 29% of the participating doctors lacked training on PPE use, 18% lacked training on COVID-19 case management, and 11% of the respondents did not receive any PPE⁴⁶. Several news reports also highlighted the logistics issues related to food, lodging and transport provision for doctors working in COVID-19 dedicated hospitals⁴⁷.

Policy recommendations

When managing any health crisis, a science-based professional response is necessary and must involve relevant experts such as public health professionals, including infectious disease epidemiologists, health policy and systems experts, medical anthropologists, health economists, and health communication experts; laboratory scientists, including virologists, microbiologists, biochemists, and lab technicians; and relevant clinicians, including physicians, nurses, and paramedics. In the long run, however, a separate public health track, which is currently absent in the Bangladeshi health sector, must be implemented³⁹. Service providers should be trained and directed to provide high-quality and efficient services with good quality and responsiveness^{48,49}, while their legitimate demands should also be duly addressed.

Research implications

Since this study did not capture the perspectives of health decision-makers, it would be beneficial to conduct further investigations into health system governance incorporating their perspectives. Quantitative research should also be conducted to explore patients' views on the responsiveness of the service providers, as well as service providers' perspectives on their own safety and experiences during the COVID-19 pandemic.

Limitations

The main limitation of this study stems from the use of online FGDs, which resulted in a sample that might not reflect the socioeconomic and demographic characteristics of the Bangladeshi population (as those without internet connectivity, or lower educational and socioeconomic status would be unable to respond to the Google Forms link or partake in online discussions). Consequently, the findings reported here cannot be generalized beyond the specific context in which the study was conducted. Second, it is worth noting that the first author was a COVID-19 patient at the time this study was conducted, which could potentially bias the qualitative analysis. However, every effort was made to reduce this risk through data triangulation⁵⁰, and by engaging multiple research team members in data coding and interpretation.

Conclusions

Bangladesh experienced several local disease outbreaks over the past several years^{51–54} as well as a dengue epidemic in 2019⁵⁵, but due to their lower magnitude compared to the COVID-19 pandemic, the need for a comprehensive overhauling of the health systems has not been felt so deeply before. Low- and middle-income countries like Bangladesh are particularly vulnerable to pandemics due to their week governance and limited health system preparedness⁵⁶. This article focused on the public perceptions of the pandemic management efforts by the health system actors, as the aim was to help the decision-makers and service providers in implementing more effective public health protection measures.

The main contribution of this investigation stems from highlighting the need to engage the right kind of experts in the right places at the outset of pandemic management efforts. It is further noted that public trust can be improved by being more transparent in official communications, while addressing the needs of service providers. These findings can help decisionmakers revise their policies in order to prevent a longer-term loss of life and economic downturn. In addressing the COVID-19 pandemic or any future public health crisis, a science-based professional response is indispensable.

Data availability

Underlying data

Harvard Dataverse: Extended data: "Public perceptions of the COVID-19 pandemic management in Bangladesh: a qualitative exploration", https://doi.org/10.7910/DVN/CJZKJH³⁴.

This project contains de-identified transcripts in Bangla.

Extended data

Harvard Dataverse: Extended data: "Public perceptions of the COVID-19 pandemic management in Bangladesh: a qualitative exploration", https://doi.org/10.7910/DVN/CJZKJH³⁴.

This project contains the following extended data:

- Discussion guide used in the FDGs in both Bangla and English.

Acknowledgments

Data are available under the terms of the Creative Commons Zero "No rights reserved" data waiver (CC0 1.0 Public domain dedication). We thank the United Nations Youth and Students Association of Bangladesh (UNYSAB) for providing logistic support for data collection. We specifically thank Professor Syed Saikh Imtiaz for his guidance and Mr. Mamun Mia for his assistance during the research activities.

References

- World Health Organization: Naming the coronavirus disease (COVID-19) and the virus that causes it. 2020; [cited 2020 21 November 2020]. Reference Source
- World Health Organization: Archived: WHO Timeline COVID-19. 2020; [cited 2020 21 November 2020]. Reference Source
- BetterWork: COVID-19 timeline in Bangladesh. 2020; [cited 2020 21 November 2020].
 Reference Source
- 4. Roser M, et al.: Coronavirus Pandemic (COVID-19). OurWorldInData.org. 2020.
- Worldometers.info: Worldometers. 2020; [cited 2020 21 November 2020]. Reference Source
- Government of Bangladesh: National Preparedness and Response Plan for COVID-19, Bangladesh. Health Services Division, Ministry of Health and Family Welfare. 2020. Reference Source
- Anwar S, Nasrullah M, Hosen MJ: COVID-19 and Bangladesh: Challenges and How to Address Them. Front Public Health. 2020; 8: 154.
 PubMed Abstract | Publisher Full Text | Free Full Text
- 8. Government of Bangladesh: Corona Info. 2020. [cited 2020 7 July 2020].
- Maswood MH: GeneXpert TB testing machines to test COVID-19. In New Age. Media New Age Limited: Dhaka. 2020. Reference Source
- Gilson L: Trust and the development of health care as a social institution. Soc Sci Med. 2003; 56(7): 1453–1468.
 PubMed Abstract | Publisher Full Text
- Ozawa S, Sripad P: How do you measure trust in the health system? A systematic review of the literature. Soc Sci Med. 2013; 91: 10–14.
 PubMed Abstract | Publisher Full Text
- Huq S, Biswas RK: COVID-19 in Bangladesh: Data deficiency to delayed decision. J Glob Health. 2020; 10(1): 010342.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Rockers PC, Kruk ME, Laugesen MJ: Perceptions of the Health System and Public Trust in Government in Low- and Middle-Income Countries: Evidence from the World Health Surveys. J Health Polit Policy Law. 2012; 37(3): 405–437.
 - PubMed Abstract | Publisher Full Text
- Department for International Development: Principles of Health Systems Resilience in the Context of Covid-19 Response. Department for International Development: UK. 2020. Reference Source
- The Joint United Nations Programme on HIV and AIDS: Rights in the time of COVID-19: Lessons from HIV for an effective, community-led response. 2020.
 Reference Source
- Kittelsen SK, Keating VC: Rational trust in resilient health systems. Health Policy Plan. 2019; 34(7): 553–557.
 PubMed Abstract | Publisher Full Text
- Roberton T, Carter ED, Chou VB, et al.: Early estimates of the indirect effects of the COVID-19 pandemic on maternal and child mortality in low-income and middle-income countries: a modelling study. Lancet Global Health. 2020; 8(7): e901-e908.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Ramalingam B, Wild L, Ferrari M: Adaptive leadership in the coronavirus response: Bridging science, policy and practice. In: Coronavirus briefing notes. ODI: London. 2020. Reference Source
- Billah MA, Miah MM, Khan MN: Reproductive number of coronavirus: A systematic review and meta-analysis based on global level evidence. *PLoS* One. 2020; 15(11): e0242128.
 PubMed Abstract | Publisher Full Text | Free Full Text

- Masrur A, Yu M, Luo W, et al.: Space-Time Patterns, Change, and Propagation of COVID-19 Risk Relative to the Intervention Scenarios in Bangladesh. Int J Environ Res Public Health. 2020; 17(16): 5911.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Khalil MM, Gain G, Mahbub-Uz-Zaman K, et al.: Gastrointestinal Manifestations among COVID-19 Patients in Bangladesh: A Cross Sectional Study. Mymensingh Med J. 2020; 29(4): 956–963. PubMed Abstract
- Khan MMA, Khan MN, Mustagir MG, et al.: Effects of underlying morbidities on the occurrence of deaths in COVID-19 patients: A systematic review and meta-analysis. J Glob Health. 2020; 10(2): 020503.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Rakib A, Sami SA, Islam MA, et al.: Epitope-Based Immunoinformatics Approach on Nucleocapsid Protein of Severe Acute Respiratory Syndrome-Coronavirus-2. Molecules. 2020; 25(21): 5088.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Abedin S, Paul SK, Nasreen SA, et al.: Molecular Detection of Human Coronavirus from North Central Part of Bangladesh Depending on ORF1ab and N Gene. Mymensingh Med J. 2020; 29(4): 865–870. PubMed Abstract
- Parvez MSA, Karim MA, Hasan M, et al.: Prediction of potential inhibitors for RNA-dependent RNA polymerase of SARS-CoV-2 using comprehensive drug repurposing and molecular docking approach. Int J Biol Macromol. 2020; 163: 1787–1797.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Do D, Sarker M, Chen S, et al.: Healthcare worker attendance during the early stages of the COVID-19 pandemic: A longitudinal analysis of fingerprint-verified data from all public-sector secondary and tertiary care facilities in Bangladesh. J Glob Health. 2020; 10(2): 020509.
 PubMed Abstract | Publisher Full Text
- Mottaleb KA, Mainuddin M, Sonobe T: COVID-19 induced economic loss and ensuring food security for vulnerable groups: Policy implications from Bangladesh. PLoS One. 2020; 15(10): e0240709.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Hamadani JD, Hasan MI, Baldi AJ, et al.: Immediate impact of stay-at-home orders to control COVID-19 transmission on socioeconomic conditions, food insecurity, mental health, and intimate partner violence in Bangladeshi women and their families: an interrupted time series. Lancet Glob Health. 2020; 8(11): e1380–e1389.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Rashid SF, Theobald S, Ozano K: Towards a socially just model: balancing hunger and response to the COVID-19 pandemic in Bangladesh. *BMJ Glob Health*. 2020; 5(6): e002715.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Ferdous MZ, Islam MS, Sikder MT, et al.: Knowledge, attitude, and practice regarding COVID-19 outbreak in Bangladesh: An online-based crosssectional study. PLoS One. 2020; 15(10): e0239254.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Hossain MA, Jahid MIK, Hossain KMA, *et al.*: Knowledge, attitudes, and fear of COVID-19 during the Rapid Rise Period in Bangladesh. *PLoS One.* 2020; 15(9): e0239646.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Paul A, Sikdar D, Hossain MM, et al.: Knowledge, attitudes, and practices toward the novel coronavirus among Bangladeshis: Implications for mitigation measures. PLoS One. 2020; 15(9): e0238492.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Bigdeli M, Rouffy B, Lane BD, et al.: Health systems governance: the missing links. BMJ Glob Health. 2020; 5(8): e002533.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Bin Khaled MN, Joardar T: Extended data: "Public perceptions of the COVID-19 pandemic management in Bangladesh: a qualitative exploration". Harvard Dataverse, V1. 2021. http://www.doi.org/10.7910/DVN/CJZKJH

- Hsieh H, Shannon S: Three approaches to qualitative content analysis. Qual Health Res. 2005; 15(9): 1277–1288.
 PubMed Abstract | Publisher Full Text
- Fake Covid-19 certificates: JKG Health Care Chairman Dr Sabrina arrested, in *The Daily Star*. Dhaka. 2020. Reference Source
- Coronavirus: Govt forms 17-member nat'l technical advisory committee, in *The Daily Star.* Dhaka. 2020. Reference Source
- Govt engages 64 secretaries to supervise relief distribution, in Prothom Alo. Bangladesh. 2020.
- Joarder T, Rawal LB, Ahmed SM, et al.: Retaining doctors in rural Bangladesh: A policy analysis. Int J Health Policy Manag. 2018; 7(9): 847–858. PubMed Abstract | Publisher Full Text | Free Full Text
- Collins T, Akselrod S, Bloomfield A, et al.: Rethinking the COVID-19 Pandemic: Back to Public Health. Ann Glob Health. 2020; 86(1): 133.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Bangladesh sees 100th death of doctors from Covid-19, in Dhaka Tribune. Dhaka. 2020.
 Reference Source
- 42. Joarder T: **How can our doctors be more responsive in the time of Covid-19 pandemic?.** In: The Business Standard. The Business Standard: Dhaka. 2020. **Reference Source**
- Joarder T, George A, Sarker M, et al.: Who are more responsive? Mixedmethods comparison of public and private sector physicians in rural Bangladesh. Health Policy Plan. 2017; 32(suppl_3): iii14–iii24.
 PubMed Abstract | Publisher Full Text
- Zaman S: Poverty and violence, frustration and inventiveness: hospital ward life in Bangladesh. Soc Sci Med. 2004; 59(10): 2025–2036.
 PubMed Abstract | Publisher Full Text
- Andaleeb SS, Siddiqui N, Khandakar S: Patient satisfaction with health services in Bangladesh. *Health Policy Plan.* 2007; 22(4): 263–273.
 PubMed Abstract | Publisher Full Text
- Islam BZ, et al.: Front Line health Workers' (FLWs) perceptions and opinions on their personal safety while attending suspected or confirmed COVID-19 patients in Bangladesh. BRAC James P Grant School of Public Health and Bangladesh Health Watch: Dhaka. 2020.

- Sakib SN: Doctors plight mounting in Bangladesh amid pandemic. In: *Andalou Agency*. Andalow Agency: Turkey. 2020. <u>Reference Source</u>
- Joarder T, George A, Ahmed SM, *et al.*: What constitutes responsiveness of physicians: A qualitative study in rural Bangladesh. *PLoS One.* 2017; 12(12): e0189962.
 - PubMed Abstract | Publisher Full Text | Free Full Text
- Joarder T, Mahmud I, Sarker M, et al.: Development and validation of a structured observation scale to measure responsiveness of physicians in rural Bangladesh. BMC Health Serv Res. 2017; 17(1): 753.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Lewis J, Lewis J, Lewis J, et al.: Generalizing from qualitative research, in Qualitative Research Practice: A Guide for Social Science Students and Researchers. J. Richie, et al.: Editors. SAGE Publications. 2013; 347.
- Maude RR, Ghose A, Samad R, et al.: A prospective study of the importance of enteric fever as a cause of non-malarial febrile illness in patients admitted to Chittagong Medical College Hospital, Bangladesh. BMC Infect Dis. 2016; 16(1): 567.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Rahim MA, Uddin KN: Chikungunya: an emerging viral infection with varied clinical presentations in Bangladesh: Reports of seven cases. *BMC Res Notes*. 2017; 10(1): 410.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Rahman M, Hossain MJ, Sultana S, et al.: Date palm sap linked to Nipah virus outbreak in Bangladesh, 2008. Vector Borne Zoonotic Dis. 2012; 12(1): 65–72. PubMed Abstract | Publisher Full Text
- Sharmin S, Viennet E, Glass K, et al.: The emergence of dengue in Bangladesh: epidemiology, challenges and future disease risk. Trans R Soc Trop Med Hyg. 2015; 109(10): 619–627.
 PubMed Abstract | Publisher Full Text
- Mutsuddy P, Jhora ST, Shamsuzzaman AKM, et al.: Dengue Situation in Bangladesh: An Epidemiological Shift in terms of Morbidity and Mortality. Can J Infect Dis Med Microbiol. 2019; 2019: 3516284.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Sriram V, Sheikh K, Soucat A: Addressing Governance Challenges and Capacities in Ministries of Health. S. Campbell, Editor. World Health Organization, UHC Partnership, Alliance for Health Policy and Systems Research: Geneva. 2020. Reference Source

Open Peer Review

Current Peer Review Status: ? ?

Version 1

Reviewer Report 26 March 2021

https://doi.org/10.5256/f1000research.31340.r80584

© **2021 Villar Uribe M.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Manuela Villar Uribe

World Bank Group, Washington, DC, USA

Thank you for inviting me to review this insightful manuscript. It comes at a very important time when discussions on lessons learned will help strengthen the health system in the short and medium term. The article is well written and presents results clearly and adequately. I only have three comments that could help strengthen the manuscript:

- 1. The participants in the focus groups are primarily students, health workers or health system experts. I suggest the results and title of the manuscript are framed more clearly to ensure the reader understands that the views presented are not actually representative of the general public but are rather the views of the health community in Bangladesh. Presenting and discussing the findings in light of the representation that the participants provide of a public health and medical population is also extremely valuable and better targeted for use in system strengthening activities to come: trust and coordination with and among medical professionals probably ought to have been a key component to the COVID response.
- 2. I am not sure I understand why you have two sections in the results where you outline issues related to decision-makers. I suggest creating a joint section for those two subsections where you can highlight the role they played and allow the reader to foresee areas of improvement at the decision-maker level.
- 3. I am particularly interested by policy recommendations and would encourage the authors to strengthen that section further by outlining recommendations in line with the results that are presented: what recommendations stem from the views on coordination, preparation, decision-makers, etc?

Is the work clearly and accurately presented and does it cite the current literature? $\ensuremath{\mathsf{Yes}}$

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others? $\ensuremath{\mathsf{Yes}}$

If applicable, is the statistical analysis and its interpretation appropriate? Not applicable

Are all the source data underlying the results available to ensure full reproducibility? $\ensuremath{\mathsf{Yes}}$

Are the conclusions drawn adequately supported by the results? Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public health, Primary Health Care, Measurement of Quality of Care, Health Policy

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Reviewer Report 24 March 2021

https://doi.org/10.5256/f1000research.31340.r80588

© **2021 Hoke T.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? Theresa Hoke

FHI 360, Durham, NC, USA

The authors have conducted a timely study investigating an intriguing aspect of the COVID-19 pandemic: public perceptions of pandemic management efforts by Bangladeshi health sector leaders. They applied sound qualitative methods to answer the research question. Three types of relationships are considered: people and decision-makers; people and providers; and providers and decision-makers.

The authors are encouraged to consider the following points as they revise the paper to make it even stronger.

Substantive issues

p.4 first column seems to indicate physicians were the only providers included, yet p.4 column 2 mentions clinicians practicing dentistry. In Bangladesh, are the latter considered physicians?

In reporting results, provide some indication of how prevalent the sentiment was across the focus

groups. For example, was the issue mentioned in just one group, or in about half the groups, or in all the groups? Similarly, if there was a strong sentiment within a group, echoed by many participants, mention that. In reading the following text, for example, it was not obvious whether this was information derived directly from the FGDs, or if it was the authors' perspective: "For years, the Bangladeshi health system has been undermined by budget shortages, lack of quality services, high out-of-pocket payments, unregulated private sector, and a highly centralized secondary or tertiary care." On p.6, "Several FGD participants commented on lack of coordination...." Were these participants in a single group, or across several groups?

On p.6: "One participant commented on the proliferation of unauthorized and even fake testing centers...." Then on p.7: "Since the COVID-19 pandemic began, a doctor has been found colluding with someone guilty of running unauthorized testing centers". Can these matters be reported together?

p.7: "Service providers' perceptions regarding health systems decision-makers include leaving them unprepared and untrained in the face of the pandemic, not recognizing their sacrifices and the lack of workplace security stemming from COVID-19 mismanagement." This text reads like a summary of findings rather than results coming directly from the FGDs. Consider moving that text to the Discussion.

p.7 The authors are encouraged to carefully consider how news articles are used to support research findings. How much confidence can be placed in the fundamental accuracy of news articles? Additionally, is it possible that news articles support what study participants say because study participants have been influenced by what they heard in the press?

Limitations: Undergraduate students represent a major portion of the non-clinician participants. This is a legitimate approach, but this particular group does bring a special perspective, since university students tend to be younger and of higher socio-economic status than the population at large. Any potential bias could be mentioned in the Limitations section.

<u>Minor issues</u>

Check wording in the following places:

Figure 1: March 26—<u>Scores</u> of people. June 16: <u>Resumption</u> of commercial...

Replace FGD's with FGDs in a couple of places.

Take another look at this wording: p.5 "For the diversity of the role among the FGD participants working in the health sector, data and observations were unique." The meaning is not completely clear.

Is the work clearly and accurately presented and does it cite the current literature? $\ensuremath{\mathsf{Yes}}$

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others? $\ensuremath{\mathsf{Yes}}$

If applicable, is the statistical analysis and its interpretation appropriate? Not applicable

Are all the source data underlying the results available to ensure full reproducibility? $\ensuremath{\mathsf{Yes}}$

Are the conclusions drawn adequately supported by the results? $\ensuremath{\mathsf{Yes}}$

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Public health; implementation research; health systems research.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000 Research