

## RESEARCH ARTICLE

# Changes in loneliness prevalence and its associated factors among Bangladeshi older adults during the COVID-19 pandemic

Sabuj Kanti Mistry<sup>1,2,3,4\*</sup>, A. R. M. Mehrab Ali<sup>1</sup>, Uday Narayan Yadav<sup>2,5,6</sup>, Fouzia Khanam<sup>7</sup>, Md. Nazmul Huda<sup>1,8,9</sup>

**1** ARCED Foundation, Dhaka, Bangladesh, **2** Centre for Primary Health Care and Equity, University of New South Wales, Sydney, Australia, **3** BRAC James P Grant School of Public Health, BRAC University, Dhaka, Bangladesh, **4** Department of Public Health, Daffodil International University, Dhaka, Bangladesh, **5** National Centre for Epidemiology and Population Health, The Australian National University, Canberra, ACT, Australia, **6** Centre for Research Policy and Implementation, Biratnagar, Nepal, **7** Department of Public Health, North South University, Dhaka, Bangladesh, **8** Translational Health Research Institute, Western Sydney University, Campeltown, NSW, Australia, **9** The School of Liberal Arts and Social Sciences, Independent University, Dhaka, Bangladesh

\* [smitra411@gmail.com](mailto:smitra411@gmail.com)



## OPEN ACCESS

**Citation:** Mistry SK, Ali ARMM, Yadav UN, Khanam F, Huda M.N (2022) Changes in loneliness prevalence and its associated factors among Bangladeshi older adults during the COVID-19 pandemic. PLoS ONE 17(11): e0277247. <https://doi.org/10.1371/journal.pone.0277247>

**Editor:** Mahfuzar Rahman, BRAC, BANGLADESH

**Received:** February 1, 2022

**Accepted:** October 23, 2022

**Published:** November 4, 2022

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0277247>

**Copyright:** © 2022 Mistry et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its [Supporting Information](#) files.

**Funding:** The author(s) received no specific funding for this work.

## Abstract

### Aims

Worldwide, loneliness is one of the most common psychological phenomena among older adults, adversely affecting their physical and mental health conditions during the COVID-19 pandemic. This study aims to assess changes in the prevalence of loneliness in the two timeframes (first and second waves of COVID-19 in Bangladesh) and identify its correlates in pooled data.

### Methods

This repeated cross-sectional study was conducted on two successive occasions (October 2020 and September 2021), overlapping with the first and second waves of the COVID-19 pandemic in Bangladesh. The survey was conducted remotely through telephone interviews among 2077 (1032 in the 2020-survey and 1045 in the 2021-survey) older Bangladeshi adults aged 60 years and above. Loneliness was measured using the 3-item UCLA Loneliness scale. The binary logistic regression model was used to identify the factors associated with loneliness in pooled data.

### Results

We found a decline in the loneliness prevalence among the participants in two survey rounds (51.5% in 2021 versus 45.7% in 2020;  $P = 0.008$ ), corresponding to 33% lower odds in the 2021-survey (AOR 0.67, 95% CI 0.54–0.84). Still, nearly half of the participants were found to be lonely in the latest survey. We also found that, compared to their respective counterparts, the odds of loneliness were significantly higher among the participants without a partner (AOR 1.58, 95% CI 1.20–2.08), with a monthly family income less than 5000 BDT (AOR 2.34, 95% CI 1.58–3.47), who lived alone (AOR 2.17, 95% CI 1.34–3.51), with poor

**Competing interests:** The authors have declared that no competing interests exist.

memory or concentration (AOR 1.58, 95% CI 1.23–2.03), and suffering from non-communicable chronic conditions (AOR 1.55, 95% CI 1.23–1.95). Various COVID-19-related characteristics, such as concern about COVID-19 (AOR 1.28, 95% CI 0.94–1.73), overwhelm by COVID-19 (AOR 1.53, 95% CI 1.14–2.06), difficulty earning (AOR 2.00, 95% CI 1.54–2.59), and receiving routine medical care during COVID-19 (AOR 2.08, 95% CI 1.61–2.68), and perception that the participants required additional care during the pandemic (AOR 2.93, 95% CI 2.27–3.79) were also associated with significantly higher odds of loneliness. However, the odds of loneliness were significantly lower among the participants with formal schooling (AOR 0.71, 95% CI 0.57–0.89) and with a family of more than four members (AOR 0.76, 95% CI 0.60–0.96).

## Conclusions

The current study found a decreased prevalence of loneliness among Bangladeshi older adults during the ongoing pandemic. However, the prevalence is still very high. The findings suggest the need for mental health interventions that may include improving social interactions increasing opportunities for meaningful social connections with family and community members and providing psychosocial support to the vulnerable population including older adults during the pandemic. It also suggests that policymakers and public health practitioners should emphasise providing mental health services at the peripheral level where the majority of older adults reside.

## Introduction

Loneliness is an emotional and mental state [1] that an individual faces in terms of subjective feelings of stress [2], sadness, low self-esteem [3], and hopelessness [4]. Loneliness is one of the most common psychological phenomena among older adults worldwide; around one-third of older adults are reported to be lonely [1, 5]. Loneliness increases the risk of heart disease, stroke, mortality, stress, chronic depression, dementia, and suicidal tendencies by damaging physical and mental health [6–9]. Empirical evidence showed that the risk of stroke and dementia raised by 30% and 50%, respectively, due to the sense of loneliness [10]. Loneliness also increased the chance of hospital visits [11], decreased quality of life [12], and mortality [13] among the older population. Additionally, one meta-analysis depicted that loneliness increases the risk of all-cause mortality by 26% among older adults [6]. Thus, loneliness in older adults remains a considerable public health concern.

The long-term COVID-19 pandemic has hard hit the world since early 2020 [14]. Recent studies showed that, amid the contemporary COVID-19 pandemic, a drastic increase in the prevalence of mental health-related issues (such as loneliness, fear, self-harm, frequent mood changes, and suicide) is seen worldwide [15–17]. Moreover, the mortality and associated physical and mental consequences of COVID-19 are disproportionately higher among the older populations [18]. The unpredictable nature of the disease, the fear of getting sick or dying, being stereotyped by others, restricted movements, home confinement for indefinite periods, limited social connectedness, and substantial and growing financial losses could aggravate the psychological situation of the general population, including older adults [19, 20].

The uncertainties and mitigating measures related to the pandemic have changed peoples' everyday lifestyles and social relationships, making them vulnerable to loneliness. The concern about loneliness among older people is particularly worrying due to their living status, need

for long-term care, and weak physical and mental health conditions [21–23]. While social distancing is essential to limit the spread of viral infection, this can negatively affect the sense of social connectedness, ultimately affecting older people's mental health. The consequence could be more severe for those with pre-existing loneliness and mental conditions [24].

Previous studies have documented that sociodemographic characteristics, such as being female, older age and unemployment, are linked to people's loneliness during the pandemic [25–29]. Poor mental and physical health, anxiety and depression can increase loneliness [25, 30, 31]. Besides, chronic conditions and functional disability have been associated with higher perceived loneliness among older adults [25, 26, 28, 29]. One study demonstrated that more worry about COVID-19 infection and more financial strain because of the pandemic was linked to greater loneliness [32]. Conversely, social resources may promote resilience among older adults during the COVID-19 pandemic [33]. Greater household size, having a partner and more emotional support are associated with less loneliness [25–27, 29, 34]. However, there are limited studies that assessed the association of loneliness with socioeconomic (e.g. income, living arrangement, distance to healthcare facility) and COVID-19 related factors, including pre-existing health conditions of COVID-19 patients, feeling concerned about the pandemic, access to medical care and financial challenges during the pandemic among older people in LMICs, including Bangladesh.

This suggests an urgent need to assess the loneliness level among older adults during the COVID-19 pandemic. To fill this need, during the first wave of the pandemic in 2020, we surveyed 1,032 older Bangladeshi adults. In 2021, the second survey was conducted among the same population to observe changes in loneliness prevalence. In the present study, we used these two cross-sectional survey data to assess changes in the prevalence of loneliness across two timeframes (first and second waves of COVID-19 in Bangladesh) and to explore the factors associated with loneliness among the Bangladeshi older population.

## Methods

### Study design and participants

This repeated cross-sectional study was conducted on two successive occasions, i.e., October 2020 and September 2021, overlapping with the first and second waves of the COVID-19 pandemic in Bangladesh. The study was conducted remotely by the Aureolin Research, Consultancy and Expertise Development (ARCED) Foundation (a non-profit organisation based in Bangladesh). The primary challenge for this study was to develop a sampling frame to select participants. Thus, we utilised our pre-established registry, constructed by merging previously completed community-based studies conducted by ARCED Foundation as described in more detail in our previous studies [35–37], which included households from all eight administrative divisions of Bangladesh, as a sampling frame. Considering the 50% prevalence of loneliness with a 5% margin of error at the 95% confidence level, 90% power of the test, and 95% response rate, a sample size of 1096 was calculated. However, during the 2020 survey, 1032 approached eligible participants responded to the study with an overall response rate of approximately 94%. During the 2021 survey, 1045 approached eligible participants responded to the study with an overall response rate of approximately 95%. Based on the population distribution of older adults by geography in Bangladesh, we adopted a probability proportionate to size (of the eight-division) approach to select older adults in each division [38]. The inclusion criterion was the minimum age of 60 years. In each administrative division, households were selected using a simple random sampling technique from the list of eligible participants in the registry. Subsequently, one eligible participant was interviewed from each of the selected households. Hence, the number of included households and respondents is equal. The exclusion criteria included adverse

mental conditions (clinically proven schizophrenia, bipolar mood disorder, dementia/cognitive impairment), a hearing disability, or an inability to communicate.

## Measures

**Outcome measure.** The study's primary outcome was loneliness, measured using the 3-item UCLA Loneliness scale [39]. The three items included: how often do you feel (i) lack of companionship, (ii) left out, and (iii) isolated in the last two weeks. Each item in the scale is measured in terms of 3-item Likert responses: hardly ever (1 point), some of the time (2 points), and often (3 points). The participants were classified as lonely if they answered 'some of the time' or 'often' to any item [1]. Dichotomised loneliness variable is used for all data analyses. We found it a reliable scale, indicated by the high internal consistency (Cronbach's alpha 0.84) among our study participants in the pooled data.

**Explanatory variables.** An extensive literature review guided the selection of explanatory variables [40–45]. Explanatory variables considered in this study were administrative division (Barishal, Chattogram, Dhaka, Mymensingh, Khulna, Rajshahi, Rangpur, Sylhet), age (categorized as 60–69, and  $\geq 70$ ), sex (male/female), marital status (married/without partner), formal schooling (without formal schooling/with formal schooling), family size ( $\leq 4$  or  $>4$ ), family monthly income (BDT) ( $<5,000$ , 5,000–10,000,  $>10,000$ ), residence (urban/rural), current occupation (employed/unemployed or retired), living arrangement (living alone or with family), walking distance to the nearest health center ( $<30$  min/ $\geq 30$  min), memory or concentration problems (no problem/low memory or concentration), suffering from non-communicable chronic conditions (yes/no), feeling concerned about COVID-19 (hardly, sometimes/often), feeling overwhelmed by COVID-19 (hardly, sometimes/often), difficulty in getting food, medicine, and routine medical care during COVID-19 (no/yes), difficulty in earning during COVID-19 (no/yes), perceived that family members are non-responsive (yes/no), and perceived that they required additional care during COVID-19 (yes/no). Self-reported information on non-communicable chronic conditions, such as arthritis, hypertension, heart diseases, stroke, hypercholesterolemia, diabetes, chronic respiratory diseases, chronic kidney disease, and cancer, was collected.

**Data collection tools and techniques.** A pre-tested semi-structured questionnaire was used to collect the information by interviewing the participants remotely using mobile phones. Data collection was accomplished electronically using SurveyCTO mobile app (<https://www.surveycpto.com/>) by trained research assistants, recruited based on previous experience administering health surveys on the electronic platform. The research assistants were trained extensively before the data collection through Zoom meetings.

The English version of the questionnaire was first translated into Bengali language and then back-translated to English by two researchers (SKM, AMA) to ensure the contents' consistency. The questionnaire was then piloted among a small sample ( $n = 10$ ) of older adults to refine the language in the final version. The tool used in the pilot study did not receive any corrections/suggestions from the participants on the contents that were developed in the local Bengali language.

**Statistical analysis.** The distribution of the variables was assessed through descriptive analyses. Given our variables' categorical nature, Chi-square tests were performed to compare differences in the prevalence of loneliness by explanatory variables, with a 5% level of significance. We used binary logistic regression models to explore the factors associated with dichotomised loneliness in the pooled data. The initial model was run with all potential covariates listed in Table 2. Then, the final model was selected using backward elimination with the Akaike information criterion (AIC). Adjusted odds ratio (aOR) and associated 95% confidence interval (95% CI) are reported in Table 3. We also performed the model diagnostics, such as

multicollinearity, the area under the curve (AUC), and the Hosmer-Lemeshow test in the model. All analyses were performed using the statistical software package Stata (Version 14.0).

## Ethical approval

The study protocol was approved by the Institutional Review Board of the Institute of Health Economics, the University of Dhaka, Bangladesh (Ref: IHE/2020/1037). Verbal informed consent was sought from the participants before administering the survey. Participation was voluntary, and participants did not receive any compensation.

## Patient and public involvement

Patients and/or the public were not involved in developing the research questions, study design, data collection and result dissemination.

## Results

### Characteristics of the participants

[Table 1](#) shows the characteristics of the study participants by survey year. In terms of survey participant coverage, there was a significant difference across geographic areas; for example, the highest coverage was from the Dhaka division in the 2020 Survey, while the highest coverage was from the Khulna division in the 2021 survey. In both surveys, most participants were 60–69 years old, male, married, without formal schooling, unemployed/retired, lived with family, and in rural areas ([Table 1](#)). However, participants' characteristics, including sex, marital status, education, and income, were significantly different across the survey years. Compared to the 2020-survey, a considerably lower proportion of participants in the 2021 survey were male (59% vs. 66%), married (77% vs. 81%), and without formal education (52% vs. 58%). The proportion of participants living with family (92% in 2020 vs. 95% in 2021), in rural areas (74% vs. 83%) and proximity to health facility (49% vs. 56%) increased significantly between the survey years. We also noted a significant increase in the reported psychological characteristics, i.e., a higher proportion of participants reported poor memory or concentration, isolation, being overwhelmed with COVID-19, and having difficulty earning and obtaining food during COVID-19 in 2021 compared to 2020 ([Table 1](#)).

### Loneliness prevalence in older adults

[Table 2](#) shows the changes in the prevalence of loneliness over time and their variation and association with participants' characteristics. We found a significant decline in the prevalence of loneliness between the two survey years (51.5% versus 45.7%;  $P = 0.008$ ). As seen in [Table 2](#), the proportion of participants experiencing loneliness decreased for almost all the characteristics presented in [Table 2](#).

Overall, the prevalence of loneliness decreased over time among females, among relatively older adults, those having a formal education, participants with higher family income, participants who were unemployed or retired, participants having memory problems, those suffering from non-communicable chronic conditions, participants who were concerned and overwhelmed by COVID-19, who faced difficulty in earning, getting medicine and routine medical care, and who perceived that they need additional care during the pandemic. However, the prevalence of loneliness seems to be static or increased a bit among the participants without a partner and those living alone ([Table 2](#)).

Table 1. Characteristics of the participants (N = 2077).

Characteristics		2020 survey		Survey 2021		P
		n	%	n	%	
Overall		1032	100.0	1045	100.0	
Administrative division						
	Barishal	149	14.4	146	14.0	0.001
	Chattogram	137	13.3	98	9.4	
	Dhaka	210	20.4	172	16.5	
	Mymensingh	63	6.1	69	6.6	
	Khulna	158	15.3	198	19.0	
	Rajshahi	103	10.0	145	13.9	
	Rangpur	144	14.0	161	15.4	
	Sylhet	68	6.6	56	5.4	
Age (year)						
	60–69	803	77.8	790	75.6	0.385
	> = 70	229	22.2	255	24.4	
Sex						
	Male	676	65.5	620	59.3	0.004
	Female	356	34.5	425	40.7	
Marital status						
	Married	840	81.4	799	76.5	0.006
	Without partner	192	18.6	246	23.5	
Formal schooling						
	Without formal schooling	602	58.3	540	51.7	0.002
	With formal schooling	430	41.7	505	48.3	
Family size						
	<4	318	30.8	347	33.2	0.243
	>4	714	69.2	698	66.8	
Family monthly income (BDT)						
	<5000	145	14.1	121	11.6	<0.001
	5000–10000	331	32.1	469	44.9	
	>10000	556	53.9	455	43.5	
Residence						
	Urban	269	26.1	182	17.4	<0.001
	Rural	763	73.9	863	82.6	
Current occupation						
	Employed	419	40.6	407	39.0	0.441
	Unemployed/retired	613	59.4	638	61.1	
Living arrangement						
	Living with family	953	92.3	992	94.9	0.016
	Living alone	79	7.7	53	5.1	
Walking distance to the nearest health centre						
	<30 minute	503	48.7	581	55.6	0.002
	≥30 minutes	529	51.3	464	44.4	
Problem in memory or concentration						
	No problem	782	75.8	676	64.7	<0.001
	Low memory or concentration	250	24.2	369	35.3	
Suffering from non communicable chronic conditions						
	No	424	41.1	447	42.8	0.435

(Continued)

**Table 1.** (Continued)

Characteristics		2020 survey		Survey 2021		P
		n	%	n	%	
Yes		608	58.9	598	57.2	
Feeling concerned about COVID-19						
	Hardly	299	29.0	348	33.3	0.033
	Sometimes to often	733	71.0	697	66.7	
Feeling overwhelmed by COVID-19						
	Hardly	370	36.4	334	32.1	0.041
	Sometimes to often	647	63.6	706	67.9	
Difficulty in getting food during COVID-19						
	No	553	55.3	514	49.7	0.011
	Yes	447	44.7	521	50.3	
Difficulty in getting medicine during COVID-19						
	No	733	75.3	764	74.8	0.765
	Yes	240	24.7	258	25.2	
Difficulty in earning during COVID-19						
	No	340	37.4	274	27.7	<0.001
	Yes	570	62.6	714	72.3	
Difficulty receiving routine medical care during COVID-19						
	No	644	69.6	709	71.0	0.517
	Yes	281	30.4	290	29.0	
Perceived that family members are non-responsive						
	No	687	66.6	738	70.6	0.047
	Yes	345	33.4	307	29.4	
Perceived needing additional care during COVID-19						
	No	769	74.5	770	73.7	0.665
	Yes	263	25.5	275	26.3	

1 USD = 85.75 BDT

<https://doi.org/10.1371/journal.pone.0277247.t001>

## Factors associated with loneliness

Table 3 shows the correlates of loneliness in the pooled sample. Compared to the 2020-survey, the odds of loneliness were significantly lower in 2021-survey (AOR 0.67, 95% CI 0.54–0.84). Compared to their respective counterparts, the odds of loneliness were significantly higher among the participants without a partner (AOR 1.58, 95% CI 1.20–2.08), living with a family having a monthly income less than 5000 BDT (AOR 2.34, 95% CI 1.58–3.47), who lived alone (AOR 2.17, 95% CI 1.34–3.51), with poor memory or concentration (AOR 1.58, 95% CI 1.23–2.03), and suffering from non-communicable chronic conditions (AOR 1.55, 95% CI 1.23–1.95). Various COVID-19-related characteristics such as concern about COVID-19 (AOR 1.28, 95% CI 0.94–1.73), overwhelmed by COVID-19 (AOR 1.53, 95% CI 1.14–2.06), difficulty earning (AOR 2.00, 95% CI 1.54–2.59), and receiving routine medical care during COVID-19 (AOR 2.08, 95% CI 1.61–2.68), the perception that they required additional care during the pandemic (AOR 2.93, 95% CI 2.27–3.79) were associated with significantly higher odds of loneliness. However, the odds of loneliness were significantly lower among the participants with formal schooling (AOR 0.71, 95% CI 0.57–0.89) and with a family of more than four members (AOR 0.76, 95% CI 0.60–0.96) (Table 3).

**Table 2.** Prevalence of loneliness over time and bivariate analysis (N = 2,077).

Characteristics	2020 survey		P <sup>1</sup>	2021 survey		P <sup>2</sup>
	n	%lonely		n	%lonely	
Overall	1032	51.5		1045	45.7	0.008
Administrative division						
Barishal	149	44.3	0.136	146	43.2	0.309
Chattogram	137	51.1		98	43.9	
Dhaka	210	57.1		172	44.2	
Mymensingh	63	46.0		69	39.1	
Khulna	158	52.5		198	46.5	
Rajshahi	103	49.5		145	41.4	
Rangpur	144	57.6		161	53.4	
Sylhet	68	42.7		56	53.6	
Age (year)						
60–69	803	49.7	0.034	790	42.8	0.001
> = 70	229	57.6		255	54.5	
Sex						
Male	676	48.5	0.009	620	40.5	<0.001
Female	356	57.0		425	53.2	
Marital status						
Married	840	50.8	0.404	799	41.2	<0.001
Without partner	192	54.2		246	60.2	
Formal schooling						
Without formal schooling	602	54.8	0.011	540	51.1	<0.001
With formal schooling	430	46.7		505	39.8	
Family size						
≤4	318	51.9	0.853	347	49.0	0.126
>4	714	51.3		698	44.0	
Family monthly income (BDT)						
<5000	145	66.2	<0.001	121	70.3	<0.001
5000–10000	331	45.0		469	43.7	
>10000	556	51.4		455	41.1	
Residence						
Urban	269	45.7	0.029	182	42.9	0.406
Rural	763	53.5		863	46.2	
Current occupation						
Employed	419	53.7	0.233	407	39.6	0.002
Unemployed/retired	613	49.9		638	49.5	
Living arrangement						
Living with family	953	49.7	<0.001	992	44.0	<0.001
Living alone	79	72.2		53	77.4	
Walking distance to the nearest health centre						
<30 minute	503	46.9	0.004	581	46.3	0.635
≥30 minutes	529	55.8		464	44.8	
Problem in memory or concentration						
No problem	782	46.4	<0.001	676	38.0	<0.001
Low memory or concentration	250	67.2		369	59.6	
Suffering from non-communicable chronic conditions						
No	424	40.8	<0.001	447	37.1	<0.001

(Continued)

**Table 2.** (Continued)

Characteristics	2020 survey			2021 survey		
	n	%lonely	P <sup>1</sup>	n	%lonely	P <sup>2</sup>
Yes	608	58.9		598	52.0	
Feeling concerned about COVID-19						
Hardly	299	31.4	<0.001	348	33.9	<0.001
Sometimes to often	733	59.6		697	51.5	
Feeling overwhelmed by COVID-19						
Hardly	370	31.9	<0.001	334	35.3	<0.001
Sometimes to often	647	62.1		706	50.4	
Difficulty in getting food during COVID-19						
No	553	38.5	<0.001	514	36.2	<0.001
Yes	447	65.3		521	54.5	
Difficulty in getting medicine during COVID-19						
No	733	44.1	<0.001	764	40.8	<0.001
Yes	240	70.4		258	58.5	
Difficulty in earning during COVID-19						
No	340	28.8	<0.001	274	29.9	<0.001
Yes	570	63.9		714	50.3	
Difficulty receiving routine medical care during COVID-19						
No	644	40.1	<0.001	709	39.4	<0.001
Yes	281	76.2		290	60.7	
Perceived that family members are non-responsive						
No	687	52.0	0.643	738	43.1	0.010
Yes	345	50.4		307	51.8	
Perceived needing additional care during COVID-19						
No	769	40.6	<0.001	770	40.0	<0.001
Yes	263	83.3		275	61.5	

P<sup>1</sup>: P-value from Pearson Chi-square test comparing participants with and without loneliness in 2020.

P<sup>2</sup>: P-value from Pearson Chi-square test comparing participants with and without loneliness in 2021.

1 USD = 85.75 BDT

<https://doi.org/10.1371/journal.pone.0277247.t002>

## Discussion

This repeated cross-sectional study assessed changes in the loneliness prevalence in older adults and its associated correlates during the first and second waves of the COVID-19 pandemic in Bangladesh. Findings found a decreased loneliness prevalence among older adults during the first and second waves of COVID-19, from 51.5% in 2020 to 45.7% in 2021.

Our study indicates that loneliness in older people during COVID-19 is still very high. To compare our study's findings, we did not find any published study that reported changes in older people's loneliness during the COVID-19 pandemic in Bangladesh and other low- and middle-income countries. However, loneliness prevalence among older adults (45.70%) in our study is broadly comparable with the rates reported by studies from high-income countries, including the United States (54%) [46], the United Kingdom (35.86%) [47], and Canada (43.1%) [18]. Meanwhile, a cross-sectional study that was conducted online on Columbian women (aged 40–59 years) found a similar rate of loneliness (44.5%) during the COVID-19 pandemic to that of the current study [48]. Pre-pandemic evidence found prevalence rates in older adults (aged ≥60 years) in Bangladesh [49] and Singapore [50], ranging from 42% to 54.3%. Many factors, including study design, study tools to measure loneliness, different time

**Table 3.** Factors associated with loneliness among the participants (N = 2077).

Characteristics		aOR	95% CI	P
Survey year				
	2020	<i>Ref</i>		
	2021	0.67	0.54–0.84	0.001
Age (year)				
	60–69	<i>Ref</i>		
	> = 70	1.18	0.90–1.54	0.237
Marital status				
	Married	<i>Ref</i>		
	Without partner	1.58	1.20–2.08	0.001
Formal schooling				
	Without formal schooling	<i>Ref</i>		
	With formal schooling	0.71	0.57–0.89	0.003
Family size				
	≤4	<i>Ref</i>		
	>4	0.76	0.60–0.96	0.021
Family monthly income (BDT)				
	>10000	<i>Ref</i>		
	5000–10000	1.01	0.80–1.29	0.907
	<5000	2.34	1.58–3.47	<0.001
Living arrangement				
	Living with family	<i>Ref</i>		
	Living alone	2.17	1.34–3.51	0.002
Walking distance to the nearest health centre				
	<30 minute	<i>Ref</i>		
	≥30 minutes	1.18	0.95–1.47	0.130
Problem in memory or concentration				
	No problem	<i>Ref</i>		
	Low memory or concentration	1.58	1.23–2.03	<0.001
Suffering from non-communicable chronic conditions				
	No	<i>Ref</i>		
	Yes	1.55	1.23–1.95	<0.001
Feeling concerned about COVID-19				
	Hardly	<i>Ref</i>		
	Sometimes to often	1.28	0.94–1.73	0.113
Feeling overwhelmed by COVID-19				
	Hardly	<i>Ref</i>		
	Sometimes to often	1.53	1.14–2.06	0.005
Difficulty in earning during COVID-19				
	No	<i>Ref</i>		
	Yes	2.00	1.54–2.59	<0.001
Difficulty receiving routine medical care during COVID-19				
	No	<i>Ref</i>		
	Yes	2.08	1.61–2.68	<0.001
Perceived that family members are non-responsive				
	No	<i>Ref</i>		
	Yes	1.25	0.98–1.59	0.068
Perceived needing additional care during COVID-19				

(Continued)

**Table 3.** (Continued)

Characteristics		aOR	95% CI	P
	No	Ref		
	Yes	2.93	2.27–3.79	<0.001

1 USD = 85.75 BDT

<https://doi.org/10.1371/journal.pone.0277247.t003>

frames, pre-and post-pandemic factors, sampling differences, cultural factors, and socio-economic contexts, may elucidate the differences in older people's loneliness during the pandemic.

The present study reported that loneliness was higher among older adults with a family with limited income, difficulty in earning and living alone. The declining traditional value system wherein family members live together and care for each other [51] and the increasing trend of nuclear families [52] are the potential reasons for older people's loneliness in Bangladesh. Our study's findings are reasonably comparable with existing literature, suggesting increased loneliness among older people with children, limited income and living alone in the community [53]. We also found a higher prevalence of loneliness in older people with no partners, those having poor memory or concentration and those suffering from non-communicable chronic conditions. Older people's loneliness was expected because they tend to live alone, lose spouses, family and friends [46, 54], and face chronic illness and experience hearing loss more than younger individuals [55]. The current study's increased loneliness in older adults may be due to COVID-19-related physical distancing and isolation measures [53]. Furthermore, limited participation in religious activities [56], limited regular activities (e.g. physical activity, exercise etc.) [57, 58], and quality of interpersonal relationships with them [48] might explain such increased loneliness in older adults. Our findings suggest loneliness prevention intervention that can address specific needs of older adults such as health conditions, housing environment, level of connectedness to close ones and cultural characteristics, the degree of loneliness experienced and the available supportive environment required for maintaining a good quality of life.

Our study found that participants feeling concerned about and overwhelmed by COVID-19, those perceiving that they required additional care, and receiving routine medical care during the pandemic were more likely to feel lonely. We could not find any related studies that had been conducted on loneliness in older adults in Bangladesh and beyond. Thus, to our knowledge, there are no or limited data available for comparison in this context. However, evidence indicates that the emergency of coronavirus interrupted [59] and overwhelmed individuals' lives, thus requiring additional care during the pandemic [44, 60]. Furthermore, people's access to public transport was limited due to coronavirus-related lockdowns [59]. As most people in Bangladesh primarily depend on public transport, such limited transportation services interrupted their routine medical care [61]. Lockdowns during the pandemic may also increase individuals' sedentary behaviours and limit physical activity [62], heightening their tension, anxiety, and fear [48] and resulting in adverse psychological health [63]. During the pandemic, such changes in individuals' everyday lives may increase the risk of loneliness in older adults [5]. Our findings suggest providing targeted care and services for reducing loneliness in older adults while maintaining Covid-19-related physical distancing measures.

To our knowledge, using a repeated cross-sectional survey, this is the first study examining the association of loneliness in older adults with their schooling and family size during the COVID-19 pandemic. As expected, decreased loneliness was found among participants with formal schooling and a family of more than four members in our study and previous research

also documented similar results [33–35]. The potential reasons are that participants with formal schooling are educated and may be aware of the adverse effects of loneliness, and the potential strategies for preventing and responding to loneliness [64]. Education can minimise individuals' loneliness by enhancing social networks and connectedness with friends and external individuals via social media [65]. Furthermore, older people with more family members have the potential to mix and interact with many family members, which reduces their risk of being lonely.

### Strengths and limitations of the study

The current study has several strengths. First, this study is among the first in Bangladesh to estimate the changes in loneliness prevalence and examine its correlates among Bangladeshi older adults during the two timeframes (first and second waves of COVID-19 in Bangladesh). Our study contributed to the limited international literature [18, 46, 47, 66] that has examined loneliness prevalence among older adults and its predictors during the COVID-19 pandemic. Second, to the best of our knowledge, some of the correlates of loneliness in older adults (e.g. feeling overwhelmed by the COVID-19 pandemic, perceiving that they required additional care during the pandemic, and receiving routine medical care) in the current study have been reported for the first time in Bangladesh and globally. Despite these strengths, our study's findings have several limitations. First, our research was cross-sectional in nature. Therefore, causality cannot be established. Second, our study is limited to quantitative analysis, as we did not explore the qualitative aspects of older adults' feelings of loneliness during the first and second waves of the pandemic. Secondly, amidst the pandemic, the sampling frame for the study was constructed by merging the contact information of previously completed community-based studies conducted by ARCED Foundation. Also, we had to conduct telephone interviews, and the sample might likely exclude those who do not have telephone access. Therefore, our sample may not represent the entire older adults of Bangladesh. These limitations highlight the need for future nationally representative research with a mixed-method approach, including a qualitative study exploring older adults' experience of loneliness and its associated factors during the COVID-19 pandemic. This will provide a better understanding of older adults' feelings of loneliness and the related factors during the COVID-19 pandemic in Bangladesh.

### Implications for policy and practice

Our findings have several implications for future research and policy for older people in Bangladesh. Firstly, our study's findings highlight the necessity for interventions to engage older adults in social and community activities and improve social interactions and community participation (while practising safety measures to limit the COVID-19 spread) to minimise the likelihood of experiencing loneliness. Secondly, the current study emphasises that it is vital to reduce the higher loneliness prevalence among older people with limited income, difficulty earning and living alone by providing financial support. Older people's loneliness can also be reduced by encouraging their children to contact and look after their parents and provide them with food and shelter. In this context, implementing the existing Parents Care Act (2013) [67] may help older people, including those who are poor, live alone, have earning difficulty, and integrate into their families. Such integration may help older people interact and live with their family members, including children and grandchildren, thus reducing their loneliness. Thirdly, it is equally important to change lonely older people's perceptions and feelings of disconnection during the pandemic. Disseminating loneliness prevention strategies via information sessions and leaflets might be a potential option for minimising older people's loneliness during the pandemic.

## Conclusion

The present study revealed that while the prevalence of loneliness decreased among Bangladeshi older adults during the pandemic, still nearly half of the participants were lonely, which needs to be taken seriously. The study's findings suggest the need for designing and implementing people-centered supportive mental health interventions for older adults to improve social interactions increasing opportunities for meaningful social connections with family and community members during this pandemic and beyond. While designing interventions, addressing the factors associated with loneliness is crucial. Policymakers and health care practitioners should also consider strengthening social support care and providing psychosocial support for the older population as part of the emergency management plan during this COVID-19 pandemic.

## Supporting information

### S1 Data.

(DTA)

## Acknowledgments

We acknowledge the role of Sadia Sumaia Chowdhury, Programme Manager, ARCED Foundation and Md. Zahirul Islam, Project Associate, ARCED Foundation, for their support in data collection for the study.

## Author Contributions

**Conceptualization:** Sabuj Kanti Mistry, A. R. M. Mehrab Ali, Uday Narayan Yadav.

**Data curation:** Sabuj Kanti Mistry.

**Formal analysis:** Sabuj Kanti Mistry.

**Investigation:** Sabuj Kanti Mistry.

**Methodology:** Sabuj Kanti Mistry.

**Project administration:** Sabuj Kanti Mistry, A. R. M. Mehrab Ali.

**Supervision:** Sabuj Kanti Mistry, A. R. M. Mehrab Ali.

**Validation:** Sabuj Kanti Mistry.

**Writing – original draft:** Sabuj Kanti Mistry, A. R. M. Mehrab Ali, Uday Narayan Yadav, Fouzia Khanam, Md. Nazmul Huda.

**Writing – review & editing:** Sabuj Kanti Mistry, Uday Narayan Yadav, Md. Nazmul Huda.

## References

1. Perissinotto CM, Cenzer IS, Covinsky KE. Loneliness in older persons: a predictor of functional decline and death. *Archives of internal medicine*. 2012; 172(14):1078–84. <https://doi.org/10.1001/archinternmed.2012.1993> PMID: 22710744
2. Burke TJ, Segrin C. Bonded or stuck? Effects of personal and constraint commitment on loneliness and stress. *Personality and Individual Differences*. 2014; 64:101–6.
3. Świtaj P, Grygiel P, Ańczewska M, Wciórka J. Experiences of discrimination and the feelings of loneliness in people with psychotic disorders: the mediating effects of self-esteem and support seeking. *Comprehensive Psychiatry*. 2015; 59:73–9. <https://doi.org/10.1016/j.comppsych.2015.02.016> PMID: 25764908

4. Chang EC, Lian X, Yu T, Qu J, Zhang B, Jia W, et al. Loneliness under assault: Understanding the impact of sexual assault on the relation between loneliness and suicidal risk in college students. *Personality and individual differences*. 2015; 72:155–9.
5. Wu B. Social isolation and loneliness among older adults in the context of COVID-19: a global challenge. *Global health research and policy*. 2020; 5(1):1–3. <https://doi.org/10.1186/s41256-020-00154-3> PMID: 32514427
6. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social isolation as risk factors for mortality: a meta-analytic review. *Perspectives on psychological science*. 2015; 10(2):227–37. <https://doi.org/10.1177/1745691614568352> PMID: 25910392
7. Hoogendoijk EO, Smit AP, van Dam C, Schuster NA, de Breij S, Holwerda TJ, et al. Frailty combined with loneliness or social isolation: an elevated risk for mortality in later life. *Journal of the American Geriatrics Society*. 2020; 68(11):2587–93. <https://doi.org/10.1111/jgs.16716> PMID: 32700319
8. Valtorta NK, Kanaan M, Gilbody S, Ronzi S, Hanratty B. Loneliness and social isolation as risk factors for coronary heart disease and stroke: systematic review and meta-analysis of longitudinal observational studies. *Heart*. 2016; 102(13):1009–16. <https://doi.org/10.1136/heartjnl-2015-308790> PMID: 27091846
9. Cacioppo JT, Hughes ME, Waite LJ, Hawkley LC, Thisted RA. Loneliness as a specific risk factor for depressive symptoms: cross-sectional and longitudinal analyses. *Psychology and aging*. 2006; 21(1):140. <https://doi.org/10.1037/0882-7974.21.1.140> PMID: 16594799
10. Donovan NJ, Blazer D. Social isolation and loneliness in older adults: review and commentary of a National Academies report. *The American Journal of Geriatric Psychiatry*. 2020; 28(12):1233–44. <https://doi.org/10.1016/j.jagp.2020.08.005> PMID: 32919873
11. Gerst-Emerson K, Jayawardhana J. Loneliness as a public health issue: the impact of loneliness on health care utilization among older adults. *American journal of public health*. 2015; 105(5):1013–9. <https://doi.org/10.2105/AJPH.2014.302427> PMID: 25790413
12. Musich S, Wang SS, Hawkins K, Yeh CS. The impact of loneliness on quality of life and patient satisfaction among older, sicker adults. *Gerontology and Geriatric Medicine*. 2015; 1:2333721415582119. <https://doi.org/10.1177/2333721415582119> PMID: 28138454
13. Holwerda TJ, Van Tilburg TG, Deeg DJH, Schutter N, Van R, Dekker J, et al. Impact of loneliness and depression on mortality: results from the Longitudinal Ageing Study Amsterdam. *The British Journal of Psychiatry*. 2016; 209(2):127–34. <https://doi.org/10.1192/bjp.bp.115.168005> PMID: 27103680
14. Hui DS, Azhar EI, Madani TA, Ntoumi F, Kock R, Dar O, et al. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health—The latest 2019 novel coronavirus outbreak in Wuhan, China. *International journal of infectious diseases*. 2020; 91:264–6. <https://doi.org/10.1016/j.ijid.2020.01.009> PMID: 31953166
15. Dawel A, Shou Y, Smithson M, Cherbuin N, Banfield M, Calear AL, et al. The effect of COVID-19 on mental health and wellbeing in a representative sample of Australian adults. *Frontiers in psychiatry*. 2020; 11:1026.
16. Liu X, Zhu M, Zhang R, Zhang J, Zhang C, Liu P, et al. Public mental health problems during COVID-19 pandemic: a large-scale meta-analysis of the evidence. *Translational psychiatry*. 2021; 11(1):1–10.
17. Terry PC, Parsons-Smith RL, Terry VR. Mood responses associated with COVID-19 restrictions. *Frontiers in Psychology*. 2020; 11:3090. <https://doi.org/10.3389/fpsyg.2020.589598> PMID: 33312153
18. Savage RD, Wu W, Li J, Lawson A, Bronskill SE, Chamberlain SA, et al. Loneliness among older adults in the community during COVID-19: a cross-sectional survey in Canada. *BMJ open*. 2021; 11(4):e044517. <https://doi.org/10.1136/bmjopen-2020-044517> PMID: 33811054
19. Salari N, Hosseiniyan-Far A, Jalali R, Vaisi-Raygani A, Rasoulpoor S, Mohammadi M, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Globalization and health*. 2020; 16(1):1–11.
20. Polšek D, Huremović D, editor. *Psychiatry of Pandemics: a Mental Health Response to Infection Outbreak*. Springer International Publishing 2019; 185 pages; ISBN978-3-030-15346-5 (e-book), ISBN978-3-030-15345-8 (softcover). *Croatian Medical Journal*. 2020; 61(3):306.
21. Koyama Y, Nawa N, Yamaoka Y, Nishimura H, Sonoda S, Kuramochi J, et al. Interplay between social isolation and loneliness and chronic systemic inflammation during the COVID-19 pandemic in Japan: Results from U-CORONA study. *Brain, Behavior, and Immunity*. 2021; 94:51–9. <https://doi.org/10.1016/j.bbi.2021.03.007> PMID: 33705870
22. Smith BJ, Lim MH. How the COVID-19 pandemic is focusing attention on loneliness and social isolation. *Public Health Res Pract*. 2020; 30(2):3022008. <https://doi.org/10.17061/phrp3022008> PMID: 32601651

23. Leigh-Hunt N, Bagguley D, Bash K, Turner V, Turnbull S, Valtorta N, et al. An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public health*. 2017; 152:157–71. <https://doi.org/10.1016/j.puhe.2017.07.035> PMID: 28915435
24. Khan MSR, Kadoya Y. Loneliness during the COVID-19 pandemic: a comparison between older and younger people. *International journal of environmental research and public health*. 2021; 18(15):7871. <https://doi.org/10.3390/ijerph18157871> PMID: 34360164
25. Cohen-Mansfield J, Hazan H, Lerman Y, Shalom V. Correlates and predictors of loneliness in older-adults: a review of quantitative results informed by qualitative insights. *International psychogeriatrics*. 2016; 28(4):557–76. <https://doi.org/10.1017/S1041610215001532> PMID: 26424033
26. Luhmann M, Hawkley LC. Age differences in loneliness from late adolescence to oldest old age. *Developmental psychology*. 2016; 52(6):943. <https://doi.org/10.1037/dev0000117> PMID: 27148782
27. Vozikaki M, Papadaki A, Linardakis M, Philalithis A. Loneliness among older European adults: Results from the survey of health, aging and retirement in Europe. *Journal of Public Health*. 2018; 26(6):613–24.
28. Kuwert P, Knaevelsrud C, Pietrzak RH. Loneliness among older veterans in the United States: Results from the National Health and Resilience in Veterans Study. *The American Journal of Geriatric Psychiatry*. 2014; 22(6):564–9. <https://doi.org/10.1016/j.jagp.2013.02.013> PMID: 23806682
29. von Soest T, Luhmann M, Hansen T, Gerstorf D. Development of loneliness in midlife and old age: Its nature and correlates. *Journal of Personality and Social Psychology*. 2020; 118(2):388. <https://doi.org/10.1037/pspp0000219> PMID: 30284871
30. Barnett MD, Moore JM, Archuleta WP. A loneliness model of hypochondriasis among older adults: The mediating role of intolerance of uncertainty and anxious symptoms. *Archives of gerontology and geriatrics*. 2019; 83:86–90. <https://doi.org/10.1016/j.archger.2019.03.027> PMID: 30974401
31. Beutel ME, Klein EM, Brähler E, Reiner I, Jünger C, Michal M, et al. Loneliness in the general population: prevalence, determinants and relations to mental health. *BMC psychiatry*. 2017; 17(1):97. <https://doi.org/10.1186/s12888-017-1262-x> PMID: 28320380
32. Polenick CA, Perbix EA, Salvi SM, Maust DT, Birditt KS, Brooks JM. Loneliness During the COVID-19 Pandemic Among Older Adults With Chronic Conditions. *Journal of Applied Gerontology*. 2021; 0733464821996527.
33. Van Orden KA, Bower E, Lutz J, Silva C, Gallegos AM, Podgorski CA, et al. Strategies to promote social connections among older adults during “social distancing” restrictions. *The American Journal of Geriatric Psychiatry*. 2021; 29(8):816–27. <https://doi.org/10.1016/j.jagp.2020.05.004> PMID: 32425473
34. Hawkley LC, Wrblewski K, Kaiser T, Luhmann M, Schumm LP. Are US older adults getting lonelier? Age, period, and cohort differences. *Psychology and Aging*. 2019; 34(8):1144.
35. Mistry SK, Ali ARM, Akther F, Yadav UN, Harris MF. Exploring fear of COVID-19 and its correlates among older adults in Bangladesh. *Globalization and Health*. 2021; 17(1):1–9.
36. Mistry SK, Ali ARM, Hossain M, Yadav UN, Ghimire S, Rahman M, et al. Exploring depressive symptoms and its associates among Bangladeshi older adults amid COVID-19 pandemic: findings from a cross-sectional study. *Social psychiatry and psychiatric epidemiology*. 2021; 56(8):1487–97. <https://doi.org/10.1007/s00127-021-02052-6> PMID: 33661353
37. Mistry SK, Ali AM, Rahman MA, Yadav UN, Gupta B, Rahman MA, et al. Changes in tobacco use patterns during COVID-19 and their correlates among older adults in Bangladesh. *International Journal of Environmental Research and Public Health*. 2021; 18(4):1779. <https://doi.org/10.3390/ijerph18041779> PMID: 33673087
38. BBS. Elderly Population in Bangladesh: Current features and future perspectives. Dhaka: Banglaedsh Bureau of Statistics, Ministry of Bangladesh, Government of Bangladesh, 2015.
39. Hughes ME, Waite LJ, Hawkley LC, Cacioppo JT. A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Research on aging*. 2004; 26(6):655–72. <https://doi.org/10.1177/0164027504268574> PMID: 18504506
40. Abir T, Osuagwu UL, Kalimullah NA, Yazdani DMN-A, Husain T, Basak P, et al. Psychological Impact of COVID-19 Pandemic in Bangladesh: Analysis of a Cross-Sectional Survey. *Health security*. 2021; 19(5):468–78. <https://doi.org/10.1089/hs.2020.0205> PMID: 34348050
41. Mamun MA, Sakib N, Gozal D, Bhuiyan AI, Hossain S, Bodrud-Doza M, et al. The COVID-19 pandemic and serious psychological consequences in Bangladesh: a population-based nationwide study. *Journal of affective disorders*. 2021; 279:462–72. <https://doi.org/10.1016/j.jad.2020.10.036> PMID: 33120247
42. Cheruvu VK, Chiyaka ET. Prevalence of depressive symptoms among older adults who reported medical cost as a barrier to seeking health care: findings from a nationally representative sample. *BMC geriatrics*. 2019; 19(1):1–10.

43. Banna MHA, Sayeed A, Kundu S, Christopher E, Hasan MT, Begum MR, et al. The impact of the COVID-19 pandemic on the mental health of the adult population in Bangladesh: a nationwide cross-sectional study. *International Journal of Environmental Health Research.* 2020;1–12. <https://doi.org/10.1080/09603123.2020.1802409> PMID: 32741205
44. Mistry SK, Ali AM, Akther F, Yadav UN, Harris MF. Exploring fear of COVID-19 and its correlates among older adults in Bangladesh. *Globalization and Health.* 2021; 17(1):1–9.
45. Mistry SK, Ali AM, Hossain MB, Yadav UN, Ghimire S, Rahman MA, et al. Exploring depressive symptoms and its associates among Bangladeshi older adults amid COVID-19 pandemic: findings from a cross-sectional study. *Social Psychiatry and Psychiatric Epidemiology.* 2021;1–11. <https://doi.org/10.1007/s00127-021-02052-6> PMID: 33661353
46. Kotwal AA, Holt-Lunstad J, Newmark RL, Cenzer I, Smith AK, Covinsky KE, et al. Social isolation and loneliness among San Francisco Bay Area older adults during the COVID-19 shelter-in-place orders. *Journal of the American Geriatrics Society.* 2021; 69(1):20–9. <https://doi.org/10.1111/jgs.16865> PMID: 32965024
47. Li LZ, Wang S. Prevalence and predictors of general psychiatric disorders and loneliness during COVID-19 in the United Kingdom. *Psychiatry research.* 2020; 291:113267. <https://doi.org/10.1016/j.psychres.2020.113267> PMID: 32623266
48. Monterrosa-Blanco A, Monterrosa-Castro Á, González-Sequeda A. Online assessment of the perception of loneliness and associated factors in Colombian climacteric women during the COVID-19 pandemic: A cross-sectional study. *Health Promotion Perspectives.* 2021; 11(2):230–9. <https://doi.org/10.34197/hpp.2021.28> PMID: 34195047
49. Rahman MS, Rahman MA, Rahman MS. Prevalence and determinants of loneliness among older adults in Bangladesh. *International Journal of Emerging Trends in Social Sciences.* 2019; 5(2):57–64.
50. Malhotra R, Tareque MI, Saito Y, Ma S, Chiu CT, Chan A. Loneliness and health expectancy among older adults: A longitudinal population-based study. *Journal of the American Geriatrics Society.* 2021; 69(11):3092–102. <https://doi.org/10.1111/jgs.17343> PMID: 34231876
51. Hossain F. Needs of the destitute elderly of Bangladesh: A sociological inquiry. *Middle East Journal of Age & Ageing.* 2013; 10(5):21–30.
52. Barikdar A, Ahmed T, Lasker SP. The situation of the elderly in Bangladesh. *Bangladesh Journal of Bioethics.* 2016; 7(1):27–36.
53. Seifert A, Hassler B. Impact of the COVID-19 pandemic on loneliness among older adults. *Frontiers in Sociology.* 2020; 5:87. <https://doi.org/10.3389/fsoc.2020.590935> PMID: 33869519
54. Pate A. Social isolation: Its impact on the mental health and wellbeing of older Victorians: COTA Victoria Melbourne, Australia; 2014.
55. Centers for Disease Control and Prevention. Loneliness and social isolation linked to serious health conditions 2022 [cited 2022 17 January]. Available from: <https://www.cdc.gov/aging/publications/features/lonely-older-adults.html>.
56. Frenkel-Yosef M, Maytles R, Shrira A. Loneliness and its concomitants among older adults during the COVID-19 pandemic. *International psychogeriatrics.* 2020; 32(10):1257–9.
57. Bai Z, Wang Z, Shao T, Qin X, Hu Z. Association between social capital and loneliness among older adults: a cross-sectional study in Anhui Province, China. *BMC geriatrics.* 2021; 21(1):1–10.
58. Fullana MA, Hidalgo-Mazzei D, Vieta E, Radua J. Coping behaviors associated with decreased anxiety and depressive symptoms during the COVID-19 pandemic and lockdown. *Journal of Affective Disorders.* 2020; 275:80–1. <https://doi.org/10.1016/j.jad.2020.06.027> PMID: 32658829
59. Huda MN, Uddin H, Hasan MK, Malo JS, Duong MC, Rahman MA. Examining Bangladesh's responses to COVID-19 in light of Vietnam: Lessons learned. *Global Biosecurity.* 2021; 3(1).
60. Van Jaarsveld GM. The effects of COVID-19 among the elderly population: a case for closing the digital divide. *Frontiers in psychiatry.* 2020;11.
61. Shammi M, Bodrud-Doza M, Islam ARMT, Rahman MM. COVID-19 pandemic, socioeconomic crisis and human stress in resource-limited settings: a case from Bangladesh. *Heliyon.* 2020; 6(5):e04063. <https://doi.org/10.1016/j.heliyon.2020.e04063> PMID: 32462098
62. Stockwell S, Trott M, Tully M, Shin J, Barnett Y, Butler L, et al. Changes in physical activity and sedentary behaviours from before to during the COVID-19 pandemic lockdown: a systematic review. *BMJ Open Sport & Exercise Medicine.* 2021; 7(1):e000960. <https://doi.org/10.1136/bmjsem-2020-000960> PMID: 34192010
63. Violant-Holz V, Gallego-Jiménez MG, González-González CS, Muñoz-Violant S, Rodríguez MJ, Sansano-Nadal O, et al. Psychological health and physical activity levels during the COVID-19 pandemic: a systematic review. *International journal of environmental research and public health.* 2020; 17 (24):9419. <https://doi.org/10.3390/ijerph17249419> PMID: 33334073

64. Burns T. OECD education and skills today 2021 [cited 2022 18 January]. Available from: <https://oecdedutoday.com/how-can-education-help-fight-loneliness/>.
65. Faizi R, El Fkihi S. Investigating the Role of Social Networks in Enhancing Students' Learning Experience: Facebook as a Case Study. International Association for Development of the Information Society. 2018.
66. O'Shea BQ, Finlay JM, Kler J, Joseph CA, Kobayashi LC. Loneliness Among US Adults Aged  $\geq 55$  Early in the COVID-19 Pandemic: Findings From the COVID-19 Coping Study. Public Health Reports. 2021;00333549211029965.
67. International Labor Organisation. Civil, commercial and family law; old-age, invalidity and survivors benefit 2013 [cited 2022 19 January]. Available from: [http://www.ilo.org/dyn/natlex/natlex4.detail?p\\_lang=en&p\\_isn=95797](http://www.ilo.org/dyn/natlex/natlex4.detail?p_lang=en&p_isn=95797).